

Department of Gastroenterology
484 Golden Autumn Way Suite 201
Bowling Green, KY 42103
Phone: (270) 781-5111 Fax: (270) 780-0475
www.gravesgilbert.com/departments/gastroenterology

Gastroparesis Diet for Delayed Stomach Emptying

Purpose

Gastroparesis is the medical term for delayed stomach emptying. During the process of digestion, the stomach must contract to empty itself of food and liquid. Normally, it contracts about three times a minute. This empties the stomach within 90-120 minutes after eating. If contractions are sluggish or less frequent, stomach emptying is delayed. This results in bothersome and sometimes serious symptoms, as well as malnutrition, because food is not being digested properly.

Gastroparesis may be caused by various conditions such as diabetes mellitus, certain disorders of the nervous system, or certain drugs. Often however, no cause can be found although a viral infection is suspected in some. Usually, the physician prescribes medication to stimulate the stomach to contract. The purpose of the gastroparesis diet is to reduce symptoms and maintain adequate fluids and nutrition. There are three steps to the diet.

STEP 1 DIET consists of liquids, which usually leave the stomach quickly by gravity alone. Liquids prevent dehydration and keep the body supplied with vital salts and minerals.

STEP 2 DIET provides additional calories by adding a small amount of dietary fat – less than 40 gm each day. For patients with gastroparesis, fatty foods and oils should be restricted, because they delay stomach emptying. However, patients at the Step 2 level are usually able to tolerate this amount.

STEP 3 DIET is designed for long-term maintenance. Fat is limited to 50 gm per day and fibrous foods are restricted, because many plant fibers cannot be digested.

Nutrition Facts

The STEP 1 Gastroparesis Diet is inadequate in all nutrients except sodium and potassium. It should not be continued for more than three days without additional nutritional support. STEP 2 and STEP 3 Gastroparesis Diets may be inadequate in Vitamins A and C and the mineral iron. A multi-vitamin supplement is usually prescribed.



Department of Gastroenterology
484 Golden Autumn Way Suite 201
Bowling Green, KY 42103
Phone: (270) 781-5111 Fax: (270) 780-0475
www.gravesgilbert.com/departments/gastroenterology

Special Considerations

- 1. Diets must be tailored to the individual patient. This is because the degree of gastroparesis may range from severe and long-standing to mild and easily corrected. Patients may also have various medical conditions to be considered. For example, diabetes patients with gastroparesis are allowed sugar-containing liquids on the Step 1 diet, because that is their only source of carbohydrate. On the Step 2 and Step 3 diets, these patients should avoid concentrated sweets. These are noted with an asterisk (*) on the food lists.
- 2. On all of the diets, liquids and food should be eaten in small, frequent meals. This helps to maintain nutrition.

Step 1 Food Groups

Group	Recommend	Avoid
Milk & milk products	none	all
Vegetables	none	all
Fruits	none	all
Breads & grains	plain saltine crackers	all others
Meat or meat substitutes	none	all
Fats & oils	none	all
Beverages	Gatorade and soft drinks (sipped slowly throughout the day)	all others
Soups	fat-free consommé and bouillon	all others

Sample Menu Step 1



Breakfast	Lunch		Dinner		
 Gatorade 1/2 cup ginger ale 1/2 cup bouillon 3/4 cup saltine crackers 6 	 Gatorade 1/2 cup Coke 1/2 cup bouillon 3/4 cup saltine crackers 6 		 Gatorade 1/2 cup Sprite 1/2 cup bouillon 3/4 cup saltine crackers 6 		
This Sample Diet Provides the Following					
Calories	790 Fat		9 gm		
Protein	11 gm Sodium		3,531 mg		
Carbohydrates	156 gm Potassium		244 mg		

Step 2 Food Groups			
Group	Recommend	Avoid	
Milk & milk products	skim milk, products made with skim milk, low-fat yogurt, low-fat cheeses	whole milk products, creams (sour, light, heavy, whipping), half & half	
Soups	fat-free consommé & bouillon, soups made from skim milk, & fat-free broths containing pasta or noodles and allowed vegetables	soups made with cream, whole milk, or broths containing fat	
Bread & grains	breads & cereals, cream of wheat, pasta, white rice, egg noodles, low-fat crackers	oatmeal, whole grain rice, cereal, bread	
Meat & meat substitutes	eggs, peanut butter (maximum 2 Tbsp/day)	beef, poultry, fish, pork products, dried beans, pea, & lentils	



Vegetables	vegetable juice (tomato, V-8); well-cooked vegetables without skins (acorn squash, beets, carrots, mushrooms, potatoes, spinach, summer squash, strained tomato sauce, yams)	all raw vegetables, cooked vegetables with skins, beans (green, wax, lima), broccoli, Brussels sprouts, cabbage, cauliflower, celery, corn, eggplant, onions, peas, peppers, pea pods, sauerkraut, turnips, water chestnuts, zucchini
Fruits	apple juice, cranberry juice, grape juice, pineapple, prune juice, canned fruits without skins (applesauce, peaches, pears)	citrus juices, all fresh and dried fruits, canned fruits with skins (apricots, cherries, blueberries, fruit cocktail, oranges, grapefruit, pineapple, plums, persimmons)
Fats & oils	any type of fat, but only in small amounts	none
Sweets & desserts*	hard candies, caramels, puddings & custards made from skim milk, frozen yogurt, fruit ice, gelatin, ice milk, jelly, honey, syrups	high-fat desserts (cakes, pies, cookies, pastries, ice cream), fruit preserves
Beverages	Gatorade*, soft drinks* (sipped slowly throughout the day)	all others, except allowed juices
*Concentrate	d sweets	

Sample Menu Step 2				
Breakfast	Lunch	Dinner		
 skim milk 1/2 cup poached egg 1 white toast slice 	 mozzarella cheese 2 oz saltine crackers 6 	 peanut butter 1 Tbsp saltine crackers 6 		



- apple juice 1/2 cup	chicken noodle soup3/4 cupGatorade 1/2 cup	 vanilla pudding 1/2 cup grape juice 1/2 cup
Morning Snack	Afternoon Snack	Evening Snack
 ginger ale 1/2 cup canned pears 1/2 cup 	 skim milk 1/2 cup cornflakes 1/2 cup sugar 2 tsp 	 frozen yogurt 1/2 cup saltine crackers 6

This Sample Diet Provides the Following					
Calories 1,343 Fat 35 gm					
Protein 52 gm Sodium 2,639 mg					
Carbohydrates 206 gm Potassium 1,411 mg					

Step 3 Food Groups			
Group Recommend Avoid			
Milk & milk products	skim milk, products made with skim milk, low-fat yogurt, low-fat cheeses	whole milk products, creams (sour, light, heavy, whipping), half & half	
Soups	fat-free consommé & bouillon, soups made from skim milk, & fat-free broths containing pasta or noodles and allowed vegetables	soups made with cream, whole milk, or broths containing fat	



	T	
Fruits	fruit juices, canned fruits without skins (applesauce, peaches, pears)	all fresh & dried fruits, canned fruits with skins (apricots, cherries, plums, blueberries, fruit cocktail, oranges, grapefruit, pineapple, persimmons)
Meat & meat substitutes	eggs, peanut butter (2 Tbsp/day), poultry, fish, lean ground beef	fibrous meats (steaks, roasts, chops), dried beans, peas, lentils
Fats & oils	any type of fat, but only in small amounts	none
Breads & grains	breads & cereals, cream of wheat, pasta, white rice, egg noodles, low-fat crackers	oatmeal, whole grain rice, cereal, bread
Vegetables	vegetable juices (tomato V-8), well-cooked vegetables without skins (acorn squash, beets, carrots, mushrooms, potatoes, spinach, summers quash, strained tomato sauce, yams)	all raw vegetables, cooked vegetables with skins: beans (green, wax, lima), broccoli, Brussels sprouts, cabbage, cauliflower, celery, corn, eggplant, onions, peas, peppers, pea pods, sauerkraut, turnips, water chestnuts, zucchini
Sweets & desserts*	hard candies, caramels, puddings & custards made from skim milk, frozen yogurt, fruit ice, gelatin, ice milk, jelly, honey, syrups	high-fat desserts (cakes, pies, cookies, pastries, ice cream), fruit preserves
Beverages	Gatorade*, soft drinks* (sipped slowly throughout the day), coffee, tea, water (note: non-caloric beverages should be limited if patient cannot	all others, except allowed juices



	maintain adequate caloric intake)	
*Concentrated sweets		

Sample Menu Step 3				
Breakfast	Lunch	Dinner		
 skim milk 1/2 cup cream of wheat 1/2 cup sugar 2 tsp orange juice 1/2 cup white toast 1 slice margarine 1 tsp jelly 1 Tbsp 	 tuna fish 2 oz low-fat mayonnaise 2 <i>Tbsp</i> white bread 2 slices canned peaches 1/2 <i>cup</i> Gatorade 1/2 cup 	 baked chicken 2 oz white rice 1/2 cup cooked beets 1/2 cup dinner roll 1 skim milk 1/2 cup margarine 2 tsp 		
Morning Snack	Afternoon Snack	Evening Snack		
low-fat yogurt1/2 cupSprite 1/2 cup	chocolate pudding1/2 cupGinger ale 1/2 cup	· ice milk 1/2 cup · pretzels 2		

This Sample Diet Provides the Following				
Calories 1,822 Fat 42 gm				
Protein 75 gm Sodium 2,234 mg				
Carbohydrates 286 gm Potassium 2,467 mg				